

On spanning tree congestion of Hamming graphs

Kyohei Kozawa*

Yota Otachi†

October 18, 2011

Abstract

We present a tight lower bound for the spanning tree congestion of Hamming graphs.

1 Preliminaries

The spanning tree congestion of graphs has been studied intensively [2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16]. In this note, we study the spanning tree congestion of Hamming graphs. We present a lower bound for the spanning tree congestion of Hamming graphs. That is, in our terminology, we show that $\text{stc}(K_n^d) \geq \frac{1}{d}(n^d - 1) \log_n d$. It is known that $\text{stc}(K_n^d) = O\left(\frac{1}{d}n^d \log_n d\right)$ [10]. Thus our lower bound is asymptotically tight.

For a graph G , we denote its vertex set and edge set by $V(G)$ and $E(G)$, respectively. For $S \subseteq V(G)$, let $G[S]$ denote the subgraph induced by S . For an edge $e \in E(G)$, we denote by $G - e$ the graph obtained from G by deleting e . Let $N_G(v)$ denote the neighborhood of $v \in V(G)$ in G ; that is, $N_G(v) = \{u \mid \{u, v\} \in E(G)\}$. We denote the degree of a vertex $v \in V(G)$ by $\deg_G(v)$, and the maximum degree of G by $\Delta(G)$; that is, $\deg_G(v) = |N_G(v)|$ and $\Delta(G) = \max_{v \in V(G)} \deg_G(v)$. A graph G is r -regular if $\deg_G(v) = r$ for every $v \in V(G)$.

For $S \subseteq V(G)$, we denote the edge set of $G[S]$ by $\iota_G(S)$, and the *boundary edge set* by $\theta_G(S)$; that is, $\iota_G(S) = \{\{u, v\} \in E(G) \mid u, v \in S\}$ and $\theta_G(S) = \{\{u, v\} \in E(G) \mid \text{exactly one of } u, v \text{ is in } S\}$. We define the function ι and θ also for a positive integer $s \leq |V(G)|$ as $\iota_G(s) = \max_{S \subseteq V(G), |S|=s} |\iota_G(S)|$ and $\theta_G(s) = \min_{S \subseteq V(G), |S|=s} |\theta_G(S)|$. Let T be a spanning tree of a connected graph G . The *congestion* of $e \in E(T)$ as $\text{cng}_G(e) = |\theta_G(L_e)|$, where L_e is the vertex set of one of the two components of $T - e$. The *congestion of T in G* , denoted by $\text{cng}_G(T)$, is the maximum congestion over all edges in T . We define the *spanning tree congestion* of G , denoted by $\text{stc}(G)$, as the minimum congestion over all spanning trees of G .

The d -dimensional Hamming graph K_n^d is the graph with vertex set $\{0, \dots, n-1\}^d$ in which two vertices are adjacent if and only if their corresponding d -dimensional vectors differ in exactly one place. It is evident that K_n^d is $d(n-1)$ -regular. The exact value of $\text{stc}(K_n^d)$ is known [6]. Also, $\text{stc}(K_2^d)$ is determined asymptotically [8].

*Electric Power Development Co., Ltd., 6-15-1, Ginza, Chuo-ku, Tokyo, 104-8165 Japan.

†Graduate School of Information Sciences, Tohoku University, Sendai 980-8579, Japan. E-mail address: otachi@dais.is.tohoku.ac.jp

2 The lower bound

Here, we present the lower bound. We need the following three lemmas.

Lemma 2.1 ([1]). *If G is r -regular and $S \subseteq V(G)$, then $|\theta_G(S)| = r|S| - 2|t_G(S)|$.*

Lemma 2.2 ([17]). *Let G be a subgraph of K_n^d . If G has s vertices and t edges, then $2t \leq (n-1)s \log_n s$.*

Lemma 2.3 ([4, 7]). *For any connected graph G , $\text{stc}(G) \geq \min_{s=\lceil (|V(G)|-1)/\Delta(G) \rceil}^{\lfloor |V(G)|/2 \rfloor} \theta(s)$.*

Theorem 2.4. $\text{stc}(K_n^d) \geq (n^d - 1) \log_n d / d$ for $n, d \geq 3$.

Proof. Since K_n^d is $d(n-1)$ -regular, Lemmas 2.1 and 2.2 imply that $\theta_{K_n^d}(s) \geq (n-1)s(d - \log_n s)$. Let $f(s) = (n-1)s(d - \log_n s)$ and $f'(s)$ be the derived function of $f(s)$. Then $f'(s) = (n-1)(d - 1/\ln n - \log_n s)$, and thus, $f(s)$ is increasing for $(n^d - 1)/(d(n-1)) \leq s \leq n^{d-1/\ln n}$ and decreasing for $n^{d-1/\ln n} \leq s \leq n^d/2$. Therefore,

$$\begin{aligned} \min_{s=\lceil (n^d-1)/(d(n-1)) \rceil}^{\lfloor n^d/2 \rfloor} f(s) &= \min \left\{ f\left(\left\lceil \frac{n^d-1}{d(n-1)} \right\rceil\right), f\left(\left\lfloor \frac{n^d}{2} \right\rfloor\right) \right\} \\ &\geq \min \left\{ f\left(\frac{n^d-1}{d(n-1)}\right), f\left(\frac{n^d}{2}\right) \right\} \\ &= \min \left\{ \frac{n^d-1}{d} \left(d - \log_n \frac{n^d-1}{d(n-1)}\right), \frac{(n-1)n^d}{2} \left(d - \log_n \frac{n^d}{2}\right) \right\} \\ &\geq \min \left\{ \frac{n^d-1}{d} \log_n d, \frac{(n-1)n^d}{2} \log_n 2 \right\}. \end{aligned}$$

Thus, by Lemma 2.3, it holds that

$$\text{stc}(K_n^d) \geq \min \left\{ \frac{n^d-1}{d} \log_n d, \frac{(n-1)n^d}{2} \log_n 2 \right\}.$$

By a simple calculation, we can see that $\frac{n^d-1}{d} \log_n d \leq \frac{(n-1)n^d}{2} \log_n 2$ for $d = 2, 3$. Since $n^d - 1 \leq (n-1)n^d$ and $(\log_n d)/d \leq (\log_n 2)/2$ for $d \geq 4$, the theorem holds. \square

References

- [1] S. L. Bezrukov, Edge isoperimetric problems on graphs, in: L. Lovász, A. Gyárfás, G. O. H. Katona, A. Recski, L. Székely (eds.), Graph Theory and Combinatorial Biology, vol. 7 of Bolyai Soc. Math. Stud., János Bolyai Math. Soc., Budapest, 1999, pp. 157–197.
- [2] H. L. Bodlaender, F. V. Fomin, P. A. Golovach, Y. Otachi, E. J. van Leeuwen, Parameterized complexity of the spanning tree congestion problem, Algorithmica.

- [3] H. L. Bodlaender, K. Kozawa, T. Matsushima, Y. Otachi, Spanning tree congestion of k -outerplanar graphs, *Discrete Math.* 311 (2011) 1040–1045.
- [4] A. Castejón, M. I. Ostrovskii, Minimum congestion spanning trees of grids and discrete toruses, *Discuss. Math. Graph Theory* 29 (2009) 511–519.
- [5] S. W. Hruska, On tree congestion of graphs, *Discrete Math.* 308 (2008) 1801–1809.
- [6] K. Kozawa, Y. Otachi, Spanning tree congestion of rook’s graphs, *Discuss. Math. Graph Theory* 31 (2011) 753–761.
- [7] K. Kozawa, Y. Otachi, K. Yamazaki, On spanning tree congestion of graphs, *Discrete Math.* 309 (2009) 4215–4224.
- [8] H.-F. Law, Spanning tree congestion of the hypercube, *Discrete Math.* 309 (2009) 6644–6648.
- [9] H.-F. Law, M. I. Ostrovskii, Spanning tree congestion: duality and isoperimetry; with an application to multipartite graphs, *Graph Theory Notes N. Y.* 58 (2010) 18–26.
- [10] H.-F. Law, M. I. Ostrovskii, Spanning tree congestion of some product graphs, *Indian J. Math.* 52 (2011) 103–111.
- [11] C. Löwenstein, D. Rautenbach, F. Regen, On spanning tree congestion, *Discrete Math.* 309 (2009) 4653–4655.
- [12] Y. Okamoto, Y. Otachi, R. Uehara, T. Uno, Hardness results and an exact exponential algorithm for the spanning tree congestion problem, in: *Proceedings of the 8th Annual Conference on Theory and Applications of Models of Computation (TAMC 2011)*, vol. 6648 of *Lecture Notes in Comput. Sci.*, pp. 452–462, Springer-Verlag, 2011.
- [13] M. I. Ostrovskii, Minimal congestion trees, *Discrete Math.* 285 (2004) 219–226.
- [14] M. I. Ostrovskii, Minimum congestion spanning trees in planar graphs, *Discrete Math.* 310 (2010) 1204–1209.
- [15] M. I. Ostrovskii, Minimum congestion spanning trees in bipartite and random graphs, *Acta Math. Sci.* 31B (2011) 634–640.
- [16] S. Simonson, A variation on the min cut linear arrangement problem, *Math. Syst. Theory* 20 (1987) 235–252.
- [17] R. Squier, B. Torrence, A. Vogt, The number of edges in a subgraph of a Hamming graph, *Appl. Math. Lett.* 14 (2001) 701–705.